



## Bisphenol A and Adiposity in an Inner-City Birth Cohort

Lori A. Hoepner, Robin M. Whyatt, Elizabeth M. Widen,  
Abeer Hassoun, Sharon E. Oberfield, Noel T. Mueller,  
Diurka Diaz, Antonia M. Calafat, Frederica P. Perera,  
and Andrew G. Rundle

<http://dx.doi.org/10.1289/EHP205>

Received: 19 March 2015

Revised: 9 October 2015

Accepted: 19 April 2016

Published: 17 May 2016

**Note to readers with disabilities:** *EHP* will provide a [508-conformant](#) version of this article upon final publication. If you require a 508-conformant version before then, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Bisphenol A and Adiposity in an Inner-City Birth Cohort**

Lori A. Hoepner<sup>1,2</sup>, Robin M. Whyatt<sup>1,2</sup>, Elizabeth M. Widen<sup>3,4,5</sup>, Abeer Hassoun<sup>6</sup>, Sharon E. Oberfield<sup>6</sup>, Noel T. Mueller<sup>7</sup>, Diurka Diaz<sup>1,2</sup>, Antonia M. Calafat<sup>8</sup>, Frederica P. Perera<sup>1,2</sup>, and Andrew G. Rundle<sup>1,2,3</sup>

<sup>1</sup>Department of Environmental Health Sciences, Mailman School of Public Health, Columbia University, 722 W. 168<sup>th</sup> St, New York, New York, USA; <sup>2</sup>Columbia Center for Children's Environmental Health, Mailman School of Public Health, Columbia University, New York, New York, USA; <sup>3</sup>Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, New York, USA; <sup>4</sup>New York Obesity Nutrition Research Center, Columbia University Medical Center, New York, New York, USA; <sup>5</sup>Institute of Human Nutrition & Department of Medicine, College of Physicians and Surgeons, Columbia University Medical Center, New York, New York, USA; <sup>6</sup>Department of Pediatrics, Division of Pediatric Endocrinology, Diabetes and Metabolism, College of Physicians and Surgeons, Columbia University Medical Center, New York, New York, USA; <sup>7</sup>Department of Epidemiology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, Maryland, USA; <sup>8</sup>National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

**Address correspondence to** Lori A. Hoepner, Department of Environmental Health Sciences, Joseph L. Mailman School of Public Health, Columbia University, 722 West 168 Street, New York, NY 10032 USA. Telephone: 00-1-646-774-5548. Fax: 00-1-646-774-7945. E-mail:

[LAH45@columbia.edu](mailto:LAH45@columbia.edu)

**Acknowledgments:** This publication was made possible by National Institutes of Health grant P01ES09600, U.S. Environmental Protection Agency grants R82702701, RD832141, RD83450901, National Institute of Environmental Health Sciences grant RC2ES018784, and the John and Wendy Neu Family Foundation. EMW and NTM supported by National Institute of Diabetes and Digestive and Kidney Diseases (T32DK091227). EMW was also supported by an unrestricted fellowship to support research in Maternal and Child Health from PepsiCo Global R+D. The authors gratefully acknowledge the technical assistance of Manori Silva, Ella Samandar, Jim Preau, Xiaoyun Ye, Ryan Hennings, Xiaoliu Zhou and Lily Jia for measuring the urinary concentrations of phthalate metabolites and bisphenol A. We are grateful to the participating families who have so generously contributed their time and effort to the study. The manuscript contents are solely the responsibility of the grantee and do not necessarily represent the official views of the US EPA. Further, the US EPA does not endorse the purchase of any commercial products or services mentioned in the publication. The findings and conclusions in this paper are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention (CDC). Use of trade names is for identification only and does not imply endorsement by the CDC, the Public Health Service, or the US Department of Health and Human Services.

**Competing financial interests:** The authors declare they have no actual or potential financial or nonfinancial conflicts of interest to disclose.

## Abstract

**Background:** Early life exposure to the endocrine disruptor bisphenol A (BPA) may contribute to development of obesity. Prospective evidence in humans on this topic is limited.

**Objectives:** We examined prenatal and early childhood BPA exposures in relation to childhood measures of adiposity in the Columbia Center for Children's Environmental Health (CCCEH) New York City birth cohort.

**Methods:** BPA concentrations were measured in prenatal (n=375) and child ages 3 (n=408) and 5 years (n=518) spot urine samples. Childhood anthropometric and bioelectrical impedance outcomes included body mass index z-scores (BMIZ) at 5 and 7 years, and fat mass index (FMI), percent body fat (%BF), and waist circumference (WC) at 7 years. Associations were evaluated using multiple linear regression with continuous and tertile BPA concentrations.

**Results:** Prenatal urinary BPA concentrations were positively associated with child age 7 FMI (beta=0.31 kg/m<sup>2</sup>, p-value=0.04, [95%CI 0.01, 0.60]), %BF (beta=0.79, p-value=0.04, [95%CI 0.03, 1.55]), and WC (beta=1.29 cm, p-value=0.01, [95%CI 0.29, 2.30]), but not BMIZ, or change in BMIZ between ages 5 and 7 years (all p-values > 0.1). FMI results were sex-specific. Child urinary BPA concentrations were not associated with child anthropometric outcomes (all p-values > 0.05).

**Conclusions:** Analyses of the CCCEH longitudinal birth cohort found associations between prenatal urinary BPA concentrations and FMI, %BF and WC. Our results suggest that prenatal BPA exposure may contribute to developmental origins of adiposity. These findings are consistent with several prior studies, raising concern about the pervasiveness of BPA.

## Introduction

Obesity in children, defined by a body mass index (BMI) greater than or equal to the 95<sup>th</sup> percentile for age and sex, is an epidemic of great concern in the United States. According to the Robert Wood Johnson Foundation, childhood obesity rates (ages 2-19 years) have more than tripled between the years 1980-2010 (RWJF 2012). Long-term risks of childhood obesity include metabolic syndrome, type 2 diabetes, cardiovascular disease and reduced adult life expectancy, with children of minority status at greater risk for becoming obese (Shaibi et al. 2007; Singh et al. 2010; Strauss and Pollack, 2001). The 2007-2008 overall prevalence of obesity and overweight was 20.3% and 17.6% for New York City (NYC) public school children, with African American and Hispanic having higher odds than white children of being obese (O.R.=1.11, 95% CI:1.07, 1.15 and O.R.=1.48, 95% CI: 1.43, 1.53, respectively) (Rundle et al. 2012). Early-life exposure to endocrine disruptors, such as bisphenol A (BPA), may be contributing to the obesity epidemic (Grün and Blumberg 2006). Endocrine disruptors may also lead to altered metabolic regulation early in life (Chevalier and Fénelichel 2014).

BPA is a known endocrine disrupting chemical and a key component in polycarbonate plastics and epoxy resins which are commonly used in consumer products resulting in significant human exposure (Calafat et al. 2008; Vandenberg et al. 2007, 2010). Exposure pathways include oral, dermal and inhalation, with dietary intake the primary route (Wilson et al. 2003; Zalko et al. 2011). Prenatal exposure may occur *in utero* or by use of catheters during delivery (Ikezuki et al. 2002; Schönfelder et al. 2002; Vandentorren et al. 2011).

Cross-sectional studies suggest that BPA exposure is associated with obesity in children (Bhandari et al. 2013; Eng et al. 2013; Li et al. 2013; Trasande et al. 2012; Wang et al. 2012; Wells et al. 2013). However, because diet is an exposure route, reverse causality cannot be ruled

out for these cross-sectional findings as obese children may have greater exposure to BPA due to higher dietary intakes. Longitudinal data on effects of prenatal BPA exposure on postnatal adiposity are limited due to dissimilarities in outcome measures, child age and geographic reference (Braun et al. 2014; Chou et al. 2011; Harley et al. 2013; Lee et al. 2013; Philippat et al. 2014; Snijder et al. 2013; Tang et al. 2013; Valvi et al. 2013). Only two other US studies investigated BPA exposure and obesity vary in their anthropometric outcomes: BMI trajectories and waist circumference (Braun et al. 2014) and BMI, waist circumference, fat mass and body fat percent (Harley et al. 2013). We hypothesized that prenatal and early childhood BPA exposure would be positively associated with childhood adiposity at ages 5 and 7 years in a NYC inner-city birth cohort.

## **Methods**

### *Study design and population*

Study subjects are participants in the prospective birth cohort of the Columbia Center for Children's Environmental Health (CCCEH) in Northern Manhattan and the South Bronx, New York. Mother-child pairs were selected if a maternal prenatal urine sample was analyzed for BPA (n=375). We additionally included children if BPA concentration in urine was measured at child age 3 (n=408) and/or 5 years (n=518). The CCCEH study design is described elsewhere (Perera et al. 2003; Whyatt et al. 2003). Briefly, we enrolled 727 women ages 18–35 years old during their third trimester of pregnancy. Women were included if they self-identified as either African American or Dominican and had resided in Northern Manhattan or the South Bronx for at least 1 year before pregnancy. Exclusion criteria included mother's report of: cigarette smoking or use of other tobacco products during pregnancy, illicit drug use, diabetes, hypertension, known HIV, or a first prenatal visit after the 20th week of gestation. Research

staff abstracted medical records of the mother and infant at delivery to ascertain prenatal medical history and birth outcomes. We contacted participants every 3 months until child age 5 and every 6 months thereafter for questionnaire follow-up, with children's biological samples collected biennially beginning at age 3 years. The CCCEH study has retained documentation regarding loss to follow-up status and reasons for missed intervals.

Data collection was completed through age 7 years in August 2013 and laboratory measurement of urinary BPA was completed for all samples. Study procedures, questionnaires and collection of biological samples were explained to each subject at enrollment and a signed consent, approved by the IRB of Columbia University Medical Center and the Centers for Disease Control and Prevention (CDC), was obtained.

#### *Urinary biomarker collection*

For mothers, urine was collected between 1999-2006 during the third trimester of pregnancy (mean gestational age: 34 weeks; SD: 3) concurrent with administration of the baseline questionnaire by trained bilingual interviewers. In children, urine was collected concurrent with follow-up questionnaires between 2001 and 2010. All samples were spot urine samples collected at varying times of day. The date, but not the exact time of collection, was available.

The total (free plus conjugated) BPA urinary concentrations (ng/mL) were measured as previously described (Calafat et al. 2008). The limit of detection (LOD) was 0.4 ng/mL. Specific gravity, as a measure of urinary dilution, was quantified at room temperature at Columbia University with a handheld refractometer (PAL 10-S, Atago).

We also measured four urinary metabolites of di-2-ethylhexyl phthalate (DEHP) at CDC: mono-2-ethylhexyl phthalate, mono-2-ethyl-5-carboxypentyl phthalate, mono-2-ethyl-5-

hydroxyhexyl phthalate, and mono-2-ethyl-5-oxohexyl phthalate (Kato et al. 2005; Silva et al. 2007). DEHP metabolite LODs ranged from 0.2 to 1.2 ng/mL. Urinary concentrations of DEHP metabolites and BPA were positively correlated ( $r$  coefficient range: 0.16, 0.34) in the CCCEH birth cohort (Hoepner et al. 2013), and in the Health Outcomes and Measures of the Environment (HOME) study (Braun et al. 2011). BPA was undetectable prenatally (6%) and at child ages 3 years (2%) and 5 years (2%). For the few BPA and DEHP metabolites results <LOD, the value of LOD/2 was substituted, consistent with prior analyses (Whyatt et al. 2003; Whyatt et al. 2009).

#### *Anthropometric measurements*

Weight prior to pregnancy was self-reported by the mother during the third trimester. Birth weight was collected from the hospital medical record. We collected body size measurements and/or body composition data for children aged 5 years ( $n=498$ ) and 7 years ( $n=511$ ). A Detecto Cardinal 750 digital scale/stadiometer (Cardinal Scale Manufacturing Company) was used to collect standing height measurements at ages 5 and 7 years until January 2010. After January 2010, we assessed standing height with the Holtain-Harpenden Wall Mounted Stadiometer Counter 602VR (Holtain Limited). Body composition values, including fat mass, were collected with the Tanita Digital Body Mass Indicator Scale BC-148 (Tanita Corporation of America) at age 7 years. Children wore light clothing and no shoes. Body mass index  $z$ -scores (BMIZ) were quantified using the SAS programs provided by the CDC (C.D.C., 2011). Fat mass index (FMI) was calculated using the algorithm  $[\text{fat mass}(\text{kg})/\text{height}(\text{m})^2]$  (VanItallie et al. 1990). Waist circumference (WC) was measured at a level midway between the lower rib margin and iliac crest using a non-stretchable measurement tape.

#### *Statistical analysis*

BPA concentrations and the molar sum of DEHP metabolite concentrations ( $\Sigma$ DEHP) were natural log-transformed for analysis to correct for their non-normal distribution. Specific gravity values were standardized by z-score transformation to stabilize model constant estimates. The standardized specific gravity concentrations were included in regression models as a covariate except for models with BPA concentration tertiles as the predictor variables. As using percentiles is a method of ordering by rank, we arithmetically adjusted BPA concentrations at each urine collection interval to calculate tertiles by first accounting for urinary dilution and then natural log-transforming using the formula: Specific gravity adjusted/log-transformed BPA =  $\text{LN}(\text{BPA}_{\text{raw}} * [(\text{mean specific gravity for population at collection interval} - 1) / (\text{individual specific gravity} - 1)])$  (Hauser et al. 2004). In other words, the prenatal BPA concentration was first specific gravity adjusted and then natural-log transformed.

For childhood BPA exposure, we calculated the mean of BPA concentrations at ages 3 and 5 years. In the absence of two childhood urinary concentrations, the single existing concentration was used (n=51).

Linear regression analyses were conducted to determine whether maternal prenatal urinary BPA concentrations predicted birth weight, BMIZ at ages 5 and 7 years and the change in BMIZ ( $\Delta$ BMIZ) from age 5 to 7 years, as well as FMI, percent body fat (%BF) and WC at age 7 years. Additionally, linear regression was used to analyze whether child urinary BPA concentrations from age 3 years or the mean of ages 3 and 5 years predict BMIZ at ages 5 and 7 years,  $\Delta$ BMIZ from age 5 to 7 years, and FMI, %BF and WC at age 7 years. To assess the representativeness of the subset analyzed, t-tests and chi-square tests were performed to compare subjects with and without prenatal and postnatal BPA concentrations and with and without anthropometric outcomes at 5 years and 7 years of age.

Maternal prenatal urinary BPA concentrations were evaluated separately from child based on prior analyses in the CCCEH birth cohort confirming prenatal BPA concentrations were not correlated with and were significantly lower than childhood BPA concentrations (Hoepner et al. 2013). However, we separately assessed as potential confounders the prenatal BPA concentrations in childhood predictor regression models and the childhood BPA concentrations in the prenatal predictor regression models and found no confounding. Furthermore, we did not find an association between prenatal BPA concentrations with maternal weight gain during pregnancy therefore we did not consider it to be a confounder variable in our analysis.

Variables were tested for in the regression models if we had previously found they were associated with BPA and if they were expected to be associated with dietary patterns and obesity (Hoepner et al. 2013, Rundle et al. 2012). Variables were included in the model if they were significantly ( $p < 0.05$ ) associated with the outcome. The following variables were controlled for in the analyses of child anthropometric measures at ages 5 and 7 years: specific gravity,  $\ln \sum \text{DEHP}$ , race/ethnicity, dichotomous maternal pre-pregnancy obesity, child sex, birth weight, and gestational age. Models including prenatal BPA concentrations also included prenatal specific gravity and  $\ln \sum \text{DEHP}$ , whereas models including childhood BPA concentrations included childhood specific gravity and  $\ln \sum \text{DEHP}$ . Additionally, we controlled for height in the analyses of outcomes: %BF at age 7 years and WC at age 7 years.

We stratified analyses by child sex in order to explore differences in associations by sex that had been found in other studies (Harley et al. 2013). Possible interaction between BPA concentration and child sex were assessed for all BMIZ and 7 year body composition outcomes.

We conducted sensitivity analyses using baseline data and logistic regression models to estimate inverse probability weighting (IPW) for successful follow-up to assess potential bias of effect estimates due to loss to follow-up and missing anthropometric data (Curtis et al. 2007; Hernan et al. 2004; Robins et al. 2000; Rundle et al. 2012; Widen et al. 2015). To estimate the weights, we included all variables in each of our final models, mother's satisfaction with living conditions, mother's years of school completed at time of pregnancy, as well as geographic information system variables using 2000 US Census block group data aggregated to 1 km radial neighborhood buffers around the home: linguistic isolation and neighborhood socioeconomic status (Rundle et al. 2012).

For all analyses, we considered results with  $p < 0.05$  to be statistically significant. Analyses were performed using IBM SPSS Statistics version 22.0 (IBM Corp., Armonk, NY, USA) and Stata version 13.0 (StataCorp, College Station, TX, USA).

## **Results**

Demographic and clinical characteristics of the 369 participants with prenatal BPA concentrations and available birth outcomes data are presented in Table 1. Table 2 shows the characteristics of the study participants grouped by BPA concentration interval and age at which anthropometric measures were collected. There was no significant difference between subjects with and without prenatal and postnatal BPA concentrations. Additionally, there was no significant difference between subjects with and without anthropometric outcomes at 5 years and 7 years of age.

### *Prenatal BPA concentration vs. child anthropometry*

Prenatal BPA concentrations were not associated with birth weight (see Supplemental Material, Table S1). Results from analyses of fully adjusted ln-transformed prenatal BPA

concentrations and child anthropometric outcomes are shown in Table 3. Overall, prenatal BPA concentrations were not significantly associated with BMIZ at ages 5 years or 7 years or  $\Delta$ BMIZ from 5 to 7 years. Prenatal BPA concentrations were positively associated with FMI at age 7 years (beta=0.31 kg/m<sup>2</sup>, p-value=0.04, [95%CI 0.01, 0.60]). Prenatal BPA concentrations were also positively associated with %BF at age 7 years (beta=0.79, p-value=0.04) and with WC at age 7 years (beta=1.29 cm, p-value=0.01).

To determine whether the FMI results were due to the fat mass or the height component of the index, we substituted the fat mass itself as the dependent variable in the model. Prenatal BPA concentrations were positively associated with fat mass regardless of the addition of height as a covariate (adjusting for height: beta=0.55 kg, p-value=0.02; without adjustment for height: beta=0.57 kg, p-value=0.03). Prenatal BPA concentrations and height were not associated. Exclusion of the prenatal ln  $\Sigma$ DEHP covariate from the model not appreciably change the associations.

#### *Prenatal tertile analysis*

Linear regression analysis using tertiles of prenatal BPA concentrations as predictor variables, with the first tertile as the reference, suggested positive linear associations with FMI, percent body weight and WC (see Table 3).

#### *Prenatal differences by child sex*

After stratifying by sex, among girls there was a positive association between prenatal BPA concentrations and 7 year FMI and WC (see Table 4). Among boys, there was no association between prenatal BPA concentrations and anthropometric outcomes. The interaction between sex and BPA concentrations was significant (see Table 4).

#### *Postnatal BPA concentration vs. child anthropometry*

Child BPA concentrations had a borderline negative association with  $\Delta$ BMIZ from 5 to 7 years (beta=-0.10, p-value=0.06, [95%CI -0.02, 0.005]). Child BPA concentrations were not associated with FMI, %BF or WC at age 7 years (see Supplemental Material, Table S2).

#### *Postnatal tertile analysis*

Linear regression analysis using tertiles of child BPA concentrations as predictor variables, with the first tertile as the reference, was consistent with the borderline negative association with  $\Delta$ BMIZ from 5 to 7 years (see Supplemental Material, Table S2). Higher exposure to BPA may be associated with lower  $\Delta$ BMIZ

#### *Postnatal differences by child sex*

When we conducted analyses stratified by sex we found a negative association between child urinary BPA concentrations and  $\Delta$ BMIZ from 5 to 7 years for girls and no associations for boys (see Supplemental Material, Table S3). Interaction terms for postnatal BPA concentrations and sex were not significant.

#### *IPW Analysis*

Weighting the data by the inverse probability of follow-up and complete anthropometric and biomarker data collection by 7 years did not appreciably change the size of the effect estimates (data not shown).

## **Discussion**

We observed a positive association between prenatal urinary BPA concentration and childhood FMI, %BF and WC at age 7 years. These results suggest prenatal BPA exposure is associated with overall body fat and central adiposity, accounting for height. However, contrary to our hypotheses, we found that maternal urinary BPA concentrations were not associated with birth weight, childhood BMIZ at ages 5 and 7 years, and  $\Delta$ BMIZ from age 5 to 7 years. The

differences in results across anthropometric outcomes may reflect differences in the construct validity of body composition versus BMI as measures of child adiposity. BMIZ alone may not be the best measure of adiposity in prepubertal children (Mueller et al. 2013). The few pediatric studies in the US that evaluated BMI versus FMI concluded that these measures are not equivalent for determining excess adiposity in children (Weber et al. 2013). This is particularly important when considering sex as the body composition components of fat mass and lean body mass differ between sexes (Weber et al. 2013). Literature on birth cohort studies of childhood obesity outcomes related to BPA exposure is limited and only one study included longitudinal analyses of prenatal BPA exposure effects on school aged boys and girls (Braun et al. 2014; Harley et al. 2013; Philippat et al. 2014; Valvi et al. 2013).

In our analysis, we found sex-specific associations with prenatal BPA exposures and FMI at age 7 years. In the Cincinnati-based HOME study, a prospective cohort composed primarily of white (67%) and African American (27%) children (n=297), prenatal urinary BPA concentrations were not associated with BMI at ages 2-5 years (Braun et al. 2014). In the aforementioned study an increase in childhood BMIZ slope per prenatal BPA tertile was observed, but sex-specific associations were weak and based on a small sample size (Braun et al. 2014). However, accelerated growth may be due to a variety of factors including fat mass and fat free mass that can only be determined with body composition techniques.

Conversely, the "INfancia y Medio Ambiente" (INMA) population-based birth cohort in Spain reported positive associations between prenatal urinary BPA concentration and body size outcomes BMI and WC at age 4 years and no sex differences (Valvi et al. 2013). Others did not identify an association of prenatal urinary BPA with weight or WC at age 3 years in a French birth cohort subset of boys only (Philippat et al. 2014). In the Center for the Health Assessment

of Mothers and Children of Salinas (CHAMACOS) birth cohort consisting primarily (98%) of Mexican American mothers and children in California, researchers found prenatal urinary BPA concentrations to be inversely related to 9 year BMIZ and %BF for girls only; concurrent urinary BPA concentrations were positively related to 9 year BMI, fat mass and WC for both sexes (Harley et al. 2013). Similar to our results, early childhood (5 year) urinary BPA concentrations in the CHAMACOS cohort were not related to late childhood (9 year) anthropometric outcomes. In addition, *in vitro* and *in vivo* experimental studies have shown positive associations between BPA exposure and adipogenesis (García-Arevalo et al. 2014; Masuno et al. 2002, 2005; Riu et al. 2011; Rubin et al. 2001; Somm et al. 2009). Our results are an important addition to the growing literature on the potential role of BPA in the developmental origin of overweight, obese and adipose deposition.

The longitudinal cohort design of this study is a major strength as prior cross-sectional studies (Bhandari et al. 2013; Li et al. 2013; Trasande et al. 2012; Wang et al. 2012) may be biased due to associations between body size and higher food intake, leading to higher exposures to BPA from food. Strengths of our study include collecting repeated urinary BPA concentrations and directly assessment of childhood anthropometric outcomes, with the addition of bioelectrical impedance body composition measures at age 7 years. The analyses found consistent results across the outcomes of FMI, %BF, and WC, although these are inter-correlated.

Another strength of our study is our ability to control for socioeconomic and additional environmental factors, including urinary phthalate concentrations measured concurrently with BPA. Additionally, our study design provides the opportunity for analysis of adiposity in understudied inner-city minority populations. Although obesity prevalence among Hispanic children

increased by 24.2% from 2003 to 2007 (Singh et al. 2010), there is little prior information on prenatal and early childhood exposure to BPA and effects on body size outcomes among US minority populations. While obesity prevalence among African American children increased 1.75% from 2003 to 2007, according to National Health and Nutrition Examination Survey (NHANES) data from 1999 to 2012, Hispanic and African American children had an obesity prevalence of 20.9% and 20.3%, respectively (Skinner and Skelton 2014). Furthermore, although the NHANES reports on BPA concentrations among different ethnic groups, it is limited to adults and children ages 6 years and older (Calafat et al. 2008). To date, no other study has evaluated the potential BPA effects on childhood body size in a birth cohort composed entirely of more than one minority group. According to the 2010 US Census, African Americans and Hispanics are the dominant minority populations in NYC (25.5% and 28.6%, respectively) (US Census Bureau, 2012). In our NYC-based birth cohort, composed of African Americans and Dominicans mother-child dyads currently being followed through adolescence, we found associations between prenatal BPA exposures and age 7 years anthropometric outcomes measures.

The use of spot urine samples is a potential limitation of our study. Prior studies of minority populations in the US found a low interclass correlation (ICC) for BPA between serial urine samples from pregnant women in Puerto Rico (ICC=0.27) (Meeker et al. 2013) and African American and Hispanic children in NYC (ICC=0.22) (Teitelbaum et al. 2008). If, as expected, the exposure misclassification because of variability of urinary BPA concentrations is non-differential in regards to the outcome, bias towards the null is expected. Thus, due to the poor reliability of the biomarkers, the findings for FMI, %BF and WC are likely to be underestimates of the true effects.

While our study was limited by the lack of dietary data during pregnancy and childhood, accurate dietary data are extremely difficult to acquire from young children given age-related development of language skills and recall ability. Dietary measures would also require quantification of BPA in food items, which was outside the scope of this study. Another possible limitation is that body composition at age 7 years was assessed with bioelectrical impedance analysis which has been validated in some populations, but has not been compared to gold-standard reference methods in a population similar to ours (Haroun et al. 2009). There are also sex differences in pubertal development that may be associated with body composition (Ahmed et al. 1999; Blum et al. 1997; Kaplowitz 2008). Earlier puberty in girls may lead to an estrogen-mediated increase in body fat (Kaplowitz 2008). Hormonal and precocious puberty data were not available for this cohort. Therefore, pubertal timing cannot be ruled out as a possible driver of the sex-specific association observed between prenatal BPA concentrations and 7 year FMI and WC. Finally, while we had missing anthropometric and biomarker data from the children in our cohort, IPW analysis suggested that loss to follow-up did not bias our results.

## **Conclusions**

In our longitudinal birth cohort, we found positive associations between prenatal urinary BPA concentrations and adiposity measures at age 7 years: FMI, %BF, and WC. The association with FMI was sex-specific for girls. Thus, future studies of environmental effects on childhood adiposity may be guided to include puberty assessment. Our findings suggest that prenatal BPA exposure may have an effect on adiposity as children age, an effect which cannot be observed by BMI-based measures alone. Our study was the first to examine and show associations between prenatal exposure to BPA and adiposity outcomes in school-aged children. As the CCCEH

cohort ages into adolescence and emerging adulthood, follow-up studies will be critical for evaluating whether the association between prenatal BPA and adiposity persists over time.

## References

- Ahmed ML, Ong KKL, Morrell DJ, Cox L, Drayyer N, Perry L et al. 1999. Longitudinal study of leptin concentrations during puberty: sex differences and relationship to changes in body composition. *J Clin Endocr Metab* 84(3):899-905.
- Bhandari R, Xiao J, Shankar A. 2013. Urinary Bisphenol A and Obesity in US Children *Am. J. Epidemiology* 177(11):1263-1270.
- Blum WF, Englaro P, Hanitsch S, Juul A, Hertal NT, Müller J et al. 1997. Plasma leptin levels in healthy children and adolescents: dependence on body mass index, body fat mass, gender, pubertal stage and testosterone. *J Clin Endocr Metab* 82:2904-2910.
- Braun JM, Kalkbrenner AE, Calafat AM, Bernert JT, Ye X, Silva MJ, et al. 2011. Variability and predictors of urinary bisphenol A concentrations during pregnancy. *Environ Health Perspect* 119(1):131-137.
- Braun JM, Lamphear BP, Calafat AM, Deria S, Khoury J, Howe CJ, et al. 2014. Early-life Bisphenol A Exposure and Child Body Mass Index: A Prospective Cohort Study. *Environ Health Perspect* 122:1239-1245.
- C.D.C. (Centers for Disease Control and Prevention). 2011. A SAS Program for the CDC Growth Charts. Atlanta, GA: Centers for Disease Control and Prevention. Available: <http://www.cdc.gov/nccdphp/dnpao/growthcharts/resources/sas.htm> [accessed 1 Oct 2012].
- Calafat AM, Ye X, Wong L, Reidy JA, Needham LL. 2008. Exposure of the US population to bisphenol A and 4-tertiary-octylphenol: 2003-2004. *Environ. Health Perspect* 116(1): 39-44.

- Chevalier N, Fénichel P. 2014. Endocrine disruptors: new players in the pathophysiology of type 2 diabetes? *Diabetes Metab pii: S1262-3636(14)00144-X*  
doi:10.1016/j.diabet.2014.09.005.
- Chou W-C, Chen J-L, Lin C-F, Chen Y-C, Shih F-C, Chuang C-Y. 2011. Biomonitoring of bisphenol A concentrations in maternal and umbilical cord blood in regard to birth outcomes and adipokine expression: a birth cohort study in Taiwan. *Environ Health* 10:94.
- Christensen KLY, Lorber M. 2014. Exposure to BPA in children - media-based and biomonitoring-based approaches. *Toxics* 2:134-157.
- Curtis LH, Hammill BG, Eisenstein EL, Kramer JM, Anstrom KJ. 2007. Using inverse probability-weighted estimators in comparative effectiveness analyses with observational databases. *Med Care* 45(10 Supl 2):S103-7.
- Eng DS, Lee JM, Gebremariam A, Meeker JD, Peterson K, Padmanabhan V. 2013. Bisphenol A and Chronic Disease Risk Factors in US Children. *Pediatrics* 132:3 e637-e645.
- García-Arevalo M, Alonso-Magdalena P, Rebelo Dos Santos J, Quesada I, Carneiro EM, Nadal A. 2014. Exposure to bisphenol-A during pregnancy partially mimics the effects of a high-fat diet altering glucose homeostasis and gene expression in adult male mice. *PLOS One* 9(6): e100214. doi: 10.1371/journal.pone.0100214.
- Grün F, Blumberg B. 2006. Environmental obesogens: organotins and endocrine disruption via nuclear receptor signaling. *Endocrinology* 147:S50-S55.
- Harley KG, Schall RA, Chevrièr J, Tyler K, Aguirre H, Bradman A, et al. 2013. Prenatal and postnatal bisphenol A exposure and body mass index in childhood in the CHAMACOS cohort. *Environ Health Perspect* 121(4):514-520.

- Haroun C, Croker H, Viner RM, Williams JE, Darch TS, Fewtrell MS, et al. 2009. Validation of BIA in obese children and adolescents and re-evaluation in a longitudinal study. *Obesity* 17:2245-2250.
- Hauser R, Meeker JD, Park S, Silva MJ, Calafat A. 2004. Temporal variability of urinary phthalate metabolite levels in men of reproductive age. *Environ Health Perspect* 112(17):1734-1740.
- Hernan MA, Hernandez-Diaz S, Robins JM. 2004. A structural approach to selection bias. *Epidemiology* 15(5):615-25.
- Hoepner LA, Whyatt RM, Just AC, Calafat AM, Perera FP, Rundle AG. 2013. Urinary concentrations of bisphenol A in an urban minority birth cohort in New York City, prenatal through age 7 years. *Environ Res*, 122:38-44.
- Ikezuki Y, Tsutsumi O, Takai Y, Kamei Y, Taketani Y. 2002. Determination of bisphenol A concentrations in human biological fluids reveals significant early prenatal exposure. *Hum Reprod* 17(11):2839-2841.
- Kaplowitz PB. 2008. Link between body fat and the timing of puberty. *Pediatrics* 121:S208-S217.
- Kato K, Silva MJ, Needham LL, Calafat AM. 2005. Determination of 16 phthalate metabolites in urine using automated sample preparation and on-line preconcentration/high-performance liquid chromatography/tandem mass spectrometry. *Anal Chem* 77(9):2985-2991.
- Lee B-E, Park H, Hong Y-C, Ha M, Kim Y, Chang N, et al. 2013. Prenatal bisphenol A and birth outcomes: MOCEH (Mothers and Children's Environmental Health) study. *Int J Hyg Environ Health* 217(2-3):328-334.

- Li D-K, Miao M, Zhou Z, Wu C, Shi H, et al. 2013. Urine Bisphenol-A Level in Relation to Obesity and Overweight in School-Age Children. *PLoS ONE* 8(6):e65399.
- Masuno H, Kidani T, Sekiya K, Sakayama K, Shiosaka T, Yamamoto H, Honda K. 2002. Bisphenol A in combination with insulin can accelerate the conversion of 3T3-L1 fibroblasts to adipocytes. *J Lipid Res* 43:676-684.
- Meeker JD, Cantonwine DE, Rivera-Gonzalez LO, Ferguson KK, Mukherjee B, Calafat AM, et al. 2013. Distribution, variability and predictors of urinary concentrations of phenols and parabens among pregnant women in Puerto Rico. *Environ Sci Technol* 47:3439-3447.
- Mueller NT, Pereira MA, Buitrago-Lopez A, Rodriguez DC, Duran AE, Ruiz AJ, et al. 2013. Adiposity indices in the prediction of insulin resistance in prepubertal Columbian children. *Public Health Nutr* 16(2):248-255.
- Perera FP, Rauh V, Tsai W, Kinney P, Camann D, Barr D, et al. 2003. Effects of transplacental exposure to environmental pollutants on birth outcomes in a multiethnic population. *Environ Health Perspect* 111(2):201-205.
- Philippat C, Botton J, Calafat AM, Ye X, Charles M, Slama R, et al. 2014. Prenatal exposure to phenols and growth in boys. *Epidemiology* 25(5):625-635.
- Riu A, Grimaldi M, le Maire A, Bey G, Phillips K, Boulahtouf A, et al. 2011. Peroxisome proliferator-activated receptor  $\gamma$  is a target for halogenated analogs of bisphenol A. *Environ Health Perspect* 119(9):1227-1232.
- Robert Wood Johnson Foundation, *F as in Fat: How Obesity Threatens America's Future*, 2012. Available: <http://www.rwjf.org/content/dam/farm/reports/reports/2012/rwjf401318> [accessed 19 Sept 2012]

- Robins JM, Hernan MA, Brumback B. 2000. Marginal structural models and causal inference in epidemiology. *Epidemiology* 11(5):550-60.
- Rubin BS, Murray MK, Damassa DA, King JC, Soto AM. 2001. Perinatal exposure to low doses of bisphenol A affects body weight, patterns of estrous cyclicity, and plasma LH levels. *Environ Health Perspect* 109(7):675-680.
- Rundle A, Richards C, Bader MDM, Schwartz-Soicher O, Lee KK, Quinn J, et al. 2012. Individual- and school-level sociodemographic predictors of obesity among New York City public school children. *Am J Epidemiol* 176(11):986-994.
- Schönfelder G, Wittfoht W, Hopp H, Talsness CE, Paul M, Chahoud I. 2002. Parent bisphenol A accumulation in the human maternal-fetal-placental unit. *Environ. Health Perspect* 110 (11):A703-A707.
- Shaibi GQ, Cruz ML, Weigensberg MJ, Toledo-Corral CM, Lane CJ, Kelly LA, et al. 2007. Adiponectin independently predicts metabolic syndrome in overweight Latino youth. *J Clin Endocr Metab* 92(5):1809-1813.
- Silva MJ, Samandar E, Preau JL, Reidy JA, Needham LL, Calafat AM. 2007. Quantification of 22 phthalate metabolites in human urine. *J Chromatogr B Analyt Technol Biomed Life Sci* 860:106-112.
- Singh GK, Siahpush M, Kogan MD. 2010. Rising social inequalities in US childhood obesity, 2003-2007. *Ann Epidemiol* 20(1):40-52.
- Skinner AC, Skelton JA. 2014. Prevalence and Trends in Obesity and Severe Obesity Among Children in the United States, 1999-2012. *JAMA Pediatr* 168(6):561-566.

- Snijder CA, Heederik D, Pierik FH, Hofman A, Jaddoe VW, Koch HM, et al. 2013. Fetal growth and prenatal exposure to bisphenol A: the generation R study. *Environ Health Perspect* 121:393-398.
- Somm E, Schwitzgebel VM, Toulotte A, Cederroth CR, Combescure C, Nef S, et al. 2009. Perinatal exposure to bisphenol A alters early adipogenesis in the rat. *Environ Health Perspect* 117(10):1549-1555.
- Strauss RS, Pollack HA. 2001. Epidemic increase in childhood weight, 1986-1998. *JAMA* 286:2845-2848.
- Tang R, Chen M, Ding G, Chen X, Han X, Zhou K, et al. 2013. Associations of prenatal exposure to phenols with birth outcomes. *Environ Pollut* 2013, 178:115-120.
- Teitelbaum, S.L., Britton, J.A., Calafat, A.M., Ye, X., Silva, M.J., Reidy, J.A., et al., 2008. Temporal variability in urinary concentrations of phthalate metabolites, phytoestrogens and phenols among minority children in the United States. *Environ Res* 106:57-269.
- Trasande L, Attina TM, Blustein J. 2012 Association between urinary bisphenol A concentration and obesity prevalence in children and adolescents. *JAMA* 308(11):1113-1121.
- U.S. Census Bureau: State and County Quick Facts. Data derived from Population Estimates, American Community Survey, Census of Population and Housing, County Business Patterns, Economic Census, Survey of Business Owners, Building Permits, Consolidated Federal Funds Report, Census of Governments, 2012.  
Available:<http://quickfacts.census.gov/qfd/states/36/3651000.html> [accessed 1 Oct 2012]
- Valvi D, Casas M, Mendez MA, Ballesteros-Gómez A, Luque N, Rubio S, Sunyer J, Vrijheid M. 2013. Prenatal bisphenol a urine concentrations and early rapid growth and overweight risk in the offspring. *Epidemiology* 24(6):791-799.

- Vandenberg LN, Chahoud I, Heindel JJ, Padmanabhan V, Paumgartten FJR, Schoenfelder G. 2010. Urinary, circulating, and tissue biomonitoring studies indicate widespread exposure to bisphenol A. *Environ Health Perspect* 118(8):1055-1070.
- Vandenberg LN, Hauser R, Marcus M, Olea N, Welshons WV. 2007. Human exposure to bisphenol A (BPA). *Reprod Toxicol* 24:139-177.
- Vandentorren S, Zenan F, Morin L, Sarter H, Bidondo M, Oleko A, et al. 2011. Bisphenol-A and phthalates contamination of urine samples by catheters in the Elfe pilot study: implications for large-scale biomonitoring studies. *Environ Res* 111 (6):761-4.
- VanItallie TB, Yang M-U, Heynsfield SB, Funk RC, Boileau RA. 1990. Height normalized indices of the body's fat-free mass and fat mass: potentially useful indicators of nutritional status. *Am J Clin Nutr* 52:953-959.
- Wang HX, Zhou Y, Tang CX, Wu JG, Chen Y, Jiang QW. 2012. Association between bisphenol A exposure and body mass index in Chinese school children: a cross-sectional study. *Environ Health* 11(1):79.
- Weber DR, Moore RH, Leonard MB, Zemel BS. 2013. Fat and lean BMI reference curves in children and adolescents and their utility in identifying excess adiposity compared with BMI and percentage body fat. *Am J Clin Nutr* 98:49-56.
- Wells EM, Jackson LW, Koontz MB. 2014. Association between bisphenol A and waist-to-height ratio among children: National Health and Nutrition Examination Survey, 2003-2010. *Ann Epidemiol* 24:165-167.
- Whyatt RW, Barr DB, Camann DE, Kinney PL, Barr JR, Andrews HF, et al. 2003. Contemporary-use pesticides in personal air samples during pregnancy and blood

samples at delivery among urban minority mothers and newborns. *Environ Health Perspect* 111(5):749-756.

Whyatt RW, Adibi JJ, Calafat AM, Camann DE, Rauh V, Bhat HK, et al. 2009. Prenatal di(2-ethylhexyl) phthalate exposure and length of gestation among an inner-city cohort. *Pediatrics* 124:e1213-1220.

Widen E, Whyatt R, Hoepner L, Mueller N, Ramirez-Carvey J, Oberfield S, et al. 2015. Gestational weight gain and obesity, adiposity and body size in African American and Dominican children in the Bronx and Northern Manhattan. *Matern Child Nutr* doi: 10.1111/mcn.12174.

**Table 1.** Characteristics of mothers and children with available birth weight and/or prenatal weight gain, and prenatal urinary BPA measures (N=369)

<i>Categorical variables: N (%)</i>	
Sex of child	
Female	201 (54.5)
Male	168 (45.5)
Race/Ethnicity	
African-American	131 (35.5)
Dominican	238 (64.5)
Foreign Born <sup>a</sup>	
US Born	162 (44)
Foreign Born	206 (56)
Maternal Pre-pregnancy Obesity (BMI >30 kg/m <sup>2</sup> ) <sup>a</sup>	
Yes	75 (20.3)
No	285 (77.2)
Parity	
Nulliparous	168 (45)
Multiparous	201 (54)
<i>Continuous variables mean (SD)</i>	
Birth weight in g	3365 (475)
Maternal Pre-pregnancy BMI <sup>a</sup> in kg/m <sup>2</sup>	25.80 (6.00)
Prenatal BPA in ng/mL	3.03 (4.16)

<sup>a</sup> # subjects missing data on each variable: Foreign Born n=1, Pre-pregnancy Obese n=9, Pre-pregnancy BMI n=9.

**Table 2.** Characteristics of subjects with urinary BPA measures and childhood anthropometric outcomes

	Prenatal BPA		Age 3 Years BPA	Age 3 and/or 5 Years BPA
	5 Year Anthropometry (N=300)	7 Year Anthropometry (N=308)	5 Year Anthropometry (N=317)	7 Year Anthropometry (N=325)
<b>Categorical variables: N (%)</b>				
Sex of Child				
Female	165 (55)	164 (53)	164 (52)	173 (53)
Male	135 (45)	144 (47)	153 (48)	152 (47)
Race/Ethnicity				
African-American	113 (38)	113 (37)	131 (41)	136 (42)
Dominican	187 (62)	195 (63)	186 (59)	189 (58)
Pre-pregnancy Obesity (BMI $\geq$ 30 kg/m <sup>2</sup> )				
Yes	64 (21)	65 (21)	71 (22)	71 (22)
Child Overweight/Obese Status				
Overweight (BMI 85 <sup>th</sup> - 95 <sup>th</sup> %ile)	49 (16)	64 (21)	52 (16)	58 (18)
Obesity (BMI >95 <sup>th</sup> %ile)	54 (18)	72 (23)	67 (21)	86 (26)
<b>Continuous variables</b>				
Mean Pre-pregnancy Weight (gms) (SD)	67.24 (17.64)	67.58 (17.69)	68.35 (18.16)	68.78 (18.12)
Mean Pre-pregnancy BMI (SD)	25.66 (6.11)	25.79 (6.14)	25.87 (6.11)	25.95 (6.13)
Mean Birth Weight (gms) (SD)	3356 (480)	3371 (487)	3381 (492)	3385 (499)
Mean BMI (SD)	16.52 (2.73)	17.93 (3.43)	16.78 (2.82)	18.19 (3.62)
Mean BMI Z-score (SD)	0.42 (1.42)	0.79 (1.15)	0.58 (1.37)	0.86 (1.15)
Median BMI Percentile (inter-quartile range)	69.47 (32.36, 92.13)	81.12 (53.38, 94.27)	74.88 (37.40, 93.11)	81.17 (54.24, 95.72)
Mean Fat Mass (kg) (SD)	--	7.19 (3.92)	--	7.47 (4.09)

Mean FMI <sup>a</sup> (SD)	--	4.51 (2.18)	--	4.67 (2.25)
Mean Percent Body Fat <sup>a</sup> (SD)	--	24.13 (6.06)	--	24.53 (6.14)
Mean Waist Circumference <sup>a</sup> (cm) (SD)	--	23.50 (8.27)	--	23.50 (8.43)
Mean Prenatal BPA (ng/mL) (SD)	3.12 (4.42)	3.06 (4.35)	--	--
Mean Child BPA (ng/mL) (SD)	--	--	8.12 (12.70)	3 year: 8.05 (12.53) 5 year: 5.35 (6.51) 3 and/or 5 year: 6.72 (7.43)

<sup>a</sup> # subjects with prenatal BPA concentration and missing data on each variable at age 7 years: fat mass index n=8, fat mass n=8, percent body fat n=8, waist circumference n=11; # subjects with mean childhood BPA concentration and missing data on each variable at age 7 years: fat mass index n=22, fat mass n=22, percent body fat n=22, waist circumference n=28

**Table 3.** Associations between prenatal urinary BPA concentrations and child anthropometric outcomes

	Beta Coefficient (95% CI)					
	Age 5 Yrs	Change From Age 5 to 7 Yrs	Age 7 Yrs			
BPA measures	BMI Z-score		FMI	Percent Body Fat	Waist Circumference (cm)	
Continuous ln-transformed BPA concentrations <sup>a,c</sup>						
Prenatal BPA	(N=300) 0.04 (-0.16, 0.24)	(N=279) 0.06 (-0.06, 0.18)	(N=308) 0.11 (-0.04, 0.26)	(N=300) 0.31* (0.01, 0.60)	(N=300) 0.79* (0.03, 1.55)	(N=297) 1.29* (0.29, 2.30)
Tertiles of specific gravity-adjusted ln-transformed prenatal BPA concentrations (in ng/mL) <sup>b,c</sup>						
<0.33	Reference	Reference	Reference	Reference	Reference	Reference
0.33-0.98	-0.29 (-0.70, 0.12)	0.16 (-0.09, 0.40)	-0.10 (-0.42, 0.21)	0.004 (-0.61, 0.40)	0.13 (-1.45, 1.71)	0.89 (-1.21, 3.00)
>0.98	-0.09 (-0.51, 0.32)	0.16 (-0.09, 0.41)	0.10 (-0.22, 0.42)	0.47 (-0.14, 1.09)	0.73 (-0.86, 2.32)	1.93 (-0.20, 4.06)

<sup>a</sup> All analyses controlled for: maternal variables: pre-pregnancy obesity, race/ethnicity, prenatal  $\Sigma$ DEHP, prenatal urinary specific gravity; child variables: sex, birth weight, gestational age.

<sup>b</sup> All analyses controlled for: maternal variables: pre-pregnancy BMI, race/ethnicity, prenatal  $\Sigma$ DEHP; child variables: sex, birth weight, gestational age.

<sup>c</sup> Additionally, height was controlled for in analyses of percent body fat and waist circumference.

\*p < 0.05 #p < 0.1

**Table 4.** Associations<sup>a</sup> between ln-transformed prenatal urinary BPA concentrations and child anthropometric outcomes stratified by sex and age 7 year interaction analysis

Prenatal BPA (ng/mL)	Beta Coefficient (95% CI)					
	Age 5 Yrs	Change From Age 5 to 7 Yrs	Age 7 Yrs			
			BMI Z-score	FMI	Percent Body Fat	Waist Circumference (cm)
Girls	(N=165) 0.02 (-0.26, 0.30)	(N=153) 0.04 (-0.14, 0.20)	(N=164) 0.12 (-0.10, 0.33)	(N=161) 0.48* (0.5, 0.91)	(N=161) 0.74 (-0.32, 1.81)	(N=160) 1.45* (0.5, 2.85)
Boys	(N=135) 0.09 (-0.20, 0.37)	(N=126) 0.03 (-0.14, 0.21)	(N=144) 0.10 (-0.12, 0.32)	(N=139) 0.06 (-0.35, 0.47)	(N=139) 0.73 (-0.40, 1.87)	(N=137) 1.07 (-0.47, 2.61)
p-value (BPA) x (Sex)	-	-	0.41	0.04*	0.51	0.32

<sup>a</sup> All analyses controlled for: maternal variables: pre-pregnancy obesity, race/ethnicity, prenatal  $\Sigma$ DEHP, prenatal urinary specific gravity; child variables: birth weight, gestational age. Additionally, height was controlled for in analyses of percent body fat and waist circumference.

\*p < 0.05 #p < 0.1